Guatemala (Ages 13-15) Global Youth Tobacco Survey (GYTS)



FACT SHEET

The Guatemala GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Guatemala could include in a comprehensive tobacco control program.

The Guatemala GYTS was a school-based survey of students in 7th, 8th, and 9th Grade conducted in 2008.

A two-stage cluster sample design was used to produce representative data for Guatemala. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 96.2%, the class response rate was 98.9%, the student response rate was 83.6%, and the overall response rate was 79.6%. A total of 3,838 students aged 13-15 participated in the Guatemala GYTS.

Prevalence

32.8% of students had ever smoked cigarettes (Boy = 39.2%, Girl = 26.4%)

16.6% currently use any tobacco product (Boy = 19.7%, Girl = 13.3%)

11.4% currently smoke cigarettes (Boy = 13.7%, Girl = 9.1%)

7.9% currently use other tobacco products (Boy = 9.5%, Girl = 6.2%)

14.8% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

27.6% think boys and 16.6% think girls who smoke have more friends 14.2% think boys and 11.2% think girls who smoke look more attractive

Access and Availability - Current Smokers

11.7% usually smoke at home

51.7% buy cigarettes in a store

80.9% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

23.1% live in homes where others smoke in their presence

40.8% are around others who smoke in places outside their home

84.6% think smoking should be banned from public places

69.5% think smoke from others is harmful to them

25.5% have one or more parents who smoke

11.5% have most or all friends who smoke

Cessation - Current Smokers

60.1% want to stop smoking

72.1% tried to stop smoking during the past year

73.9% have ever received help to stop smoking

Media and Advertising

76.6% saw anti-smoking media messages, in the past 30 days

81.0% saw pro-cigarette ads on billboards, in the past 30 days

79.0% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

9.5% have an object with a cigarette brand logo

11.8% were offered free cigarettes by a tobacco company representative

School

57.7% had been taught in class, during the past year, about the dangers of smoking

29.6% had discussed in class, during the past year, reasons why people their age smoke

54.7% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 16.6% of the students currently use any form of tobacco; 11.4% of the students currently smoke cigarettes; 7.9% of the students currently use some other form of tobacco.
- SHS exposure is high almost one-quarter of the students live in homes where others smoke, and 2 in 5 students are exposed to smoke around others outside of the home; one-quarter of the students have at least one parent who smokes.
- Seven in 10 students think smoke from others is harmful to them.
- Over 8 in 10 students think smoking in public places should be banned.
- Three in 5 current smokers want to stop smoking.
- 9.5% of students have an object with a cigarette brand logo on it.
- Three-quarters of the students saw anti-smoking media messages in the past 30 days; 8 in 10 students saw pro-cigarette ads on billboards and in newspapers or magazines in the past 30 days.